

2017

MATHEMATICS

[**Honours**]

(CBCS)

[**First Semester**]

PAPER – C2T

Full Marks : 60

Time : 3 hours

The figures in the right hand margin indicate marks

UNIT – I

(*Classical Algebra*)

1. Answer any *one* question : 2 × 1

(a) If $x + iy$ moves on the straight line $3x + 4y + 5 = 0$, then find the minimum value of $|x + iy|$. 2

(b) Solve the equation $x^5 + x^4 + x^3 + x^2 + x + 1 = 0$. 2

2. Answer any *two* questions : 5 × 2

(a) If $(1 + i \tan \alpha)^{1 + i \tan \beta}$ can have real values, then show that one of them is $(\sec \alpha)^{\sec^2 \beta}$. 5

(b) Show that the condition that the sum of two roots of the equation $x^4 + mx^2 + nx + p = 0$ be equal to the product of the other two roots is $(2p - n)^2 = (p - n)(p + m - n)^2$. 5

(c) If a_1, a_2, \dots, a_n be n real positive quantities then prove that

$$\text{A.M.} \geq \text{G.M.} \geq \text{H.M.} \quad 5$$

3. Answer any *one* question : 10 × 1

(a) (i) If $x + \frac{1}{x} = 2 \cos \alpha$, $y + \frac{1}{y} = 2 \cos \beta$,
 $z + \frac{1}{z} = 2 \cos \gamma$, and $x + y + z = 0$ then
 prove that

$$\sum \sin 4\alpha = 2 \sum \sin(\beta + \gamma)$$

$$\text{and } \sum \cos 4\alpha = 2 \sum \cos(\beta + \gamma) \quad 5$$

(ii) If the equation whose roots are squares of the roots of the cubic $x^3 - ax^2 + bx - 1 = 0$ is identical with this cubic, prove that either $a = b = 0$ or $a = b = 3$ or a, b are the roots of the equation $t^2 + t + 2 = 0$. 5

(b) (i) If a, b, c, x, y, z be all real numbers and $a^2 + b^2 + c^2 = 1, x^2 + y^2 + z^2 = 1$ then prove that $-1 \leq ax + by + cz \leq 1$.

If a_1, a_2, \dots, a_n be n positive rational numbers and $s = a_1 + a_2 + \dots + a_n$, prove that

$$\left(\frac{s}{a_1} - 1\right)^{a_1} \left(\frac{s}{a_2} - 1\right)^{a_2} \dots \left(\frac{s}{a_n} - 1\right)^{a_n} \leq (n-1)^s. \quad 2+3$$

(ii) If the equation $x^3 + px^2 + qx + r = 0$ has a root $\alpha + i\alpha$ where p, q, r and α are real, prove that $(p^2 - 2q)(q^2 - 2pr) = r^2$.

Hence solve the equation

$$x^3 - x^2 - 4x + 24 = 0. \quad 3+2$$

UNIT – II

(Sets and Integers)

4. Answer any five questions : 2 × 5

(a) Prove that intersection of two equivalence relations is also an equivalence relation. 2

(b) Prove that square of any integer is of the form $3k$ or $3k + 1$. 2

(c) Examine if the relation ρ on the set \mathbb{Z} is an equivalence relation or not

$$\rho = \{(a, b) \in \mathbb{Z} \times \mathbb{Z} : |a - b| \leq 3\}. \quad 2$$

(d) Prove that, there exists no integer in between 0 and 1. 2

(e) Let

$$P = \{n \in \mathbb{Z} : 0 \leq n \leq 5\}, Q = \{n \in \mathbb{Z} : -5 \leq n \leq 0\}$$

be two sets. Prove that cardinality of two sets are equal. 2

(5)

(f) If $s_n = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}$ then prove that

$$s_n > \frac{2n}{n+1}$$

if $n > 1$.

2

(g) If X and Y are two non-empty sets and $f : X \rightarrow Y$ be an onto mapping, then for any subsets A and B of Y , prove that

$$f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B).$$

2

(h) (i) State the Fundamental theorem of Arithmetic.

(ii) If a divides b , then prove that every divisor of a divides b .

2

5. Answer any one question : 5 × 1

(a) (i) Prove that $1^n - 3^n - 6^n + 8^n$ is divisible by $10 \forall n \in \mathbb{N}$. 2

(ii) Find integers u and v satisfying $20u + 63v = 1$. 3

(b) (i) State the division algorithm on the set of integers. 1

(ii) Find integers s and t such that

$$\gcd(341, 1643) = 341s + 1643t. \quad 2$$

(iii) Using the theory of congruence for finding the remainder when the sum $1^5 + 2^5 + 3^5 + \dots + 100^5$ is divided by 5. 2

UNIT – III

(System of Linear Equations)

6. Answer any two questions : 2×2

(a) Solve the system of equations :

$$x + 2y - z - 3w = 1$$

$$2x + 4y + 3z + w = 3$$

$$3x + 6y + 4z - 2w = 5$$

if possible.

2

(7)

(b) For what values of k the system of equations

$$x + 2y + 3z = kx$$

$$2x + y + 3z = ky$$

$$2x + 3y + z = kz$$

has a non-trivial solution. 2

(c) Determine k so that the set $\{(1, 2, 1), (k, 3, 1), (2, k, 0)\}$ is linearly dependent. 2

7. Answer any *one* question : 5 × 1

(a) Determine the conditions for which the system

$$x + y + z = 1$$

$$x + 2y - z = b$$

$$5x + 7y + az = b^2$$

admits of (i) only one solution.

(ii) no solution.

(iii) many solutions. 5

(8)

- (b) (i) Obtain the fully row reduced normal form of the matrix : 2

$$\begin{pmatrix} 0 & 0 & 1 & 2 & 1 \\ 1 & 3 & 1 & 0 & 3 \\ 2 & 6 & 4 & 2 & 8 \\ 3 & 9 & 4 & 2 & 10 \end{pmatrix}$$

- (ii) For what values of k , the planes $x - 4y + 5z = k$, $x - y + 2z = 3$, $2x + y + z = 0$ intersect in a line. 3

UNIT - IV

(*Linear Transformation and Eigenvalues*)

8. Answer any *two* questions : 2×2

- (a) Find the rank of the matrix :

$$\begin{pmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{pmatrix}$$

if two straight lines $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$ are coincident. 2

(b) Show that the rank of a skew symmetric matrix cannot be 1. 2

(c) State Cayley-Hamilton theorem and using theorem find A^{-1} , where

$$A = \begin{pmatrix} 2 & 1 \\ 3 & 5 \end{pmatrix}. \quad 2$$

9. Answer any one question : 10 × 1

(a) (i) If $A = \begin{pmatrix} 1 & -1 \\ v_2 & v_2 \\ 1 & 1 \\ v_2 & v_2 \end{pmatrix}$,

$X = (x_1, x_2)^T$ and $Y = (y_1, y_2)^T$. Verify by means of the transformation $X = AY$ that $x_1^2 + x_2^2$ is transformed to $y_1^2 + y_2^2$.

Find the dimension of the subspace \mathbb{R}^3 defined by

$$S = \{(x, y, z) \in \mathbb{R}^3 : x + 2y = z, 2x + 3z = y\}. \quad 3 + 2$$

(ii) Verify Caley-Hamilton's theorem for the matrix

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 2 & 1 \\ 2 & 3 & 2 \end{pmatrix}$$

Hence compute A^{-1} . 3 + 2

(b) (i) Find all real λ for which the rank of the matrix A in 2, where

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & -1 & \lambda \\ 5 & 7 & 1 & \lambda^2 \end{pmatrix} \quad 3$$

(ii) If X_1, X_2, \dots, X_r be r eigen vectors of an $n \times n$ matrix A corresponding to r distinct eigen values $\lambda_1, \lambda_2, \dots, \lambda_r$ respectively, then prove that X_1, X_2, \dots, X_r are linearly independent. 5

(iii) λ is an eigen value of a real skew symmetric matrix. Prove that

$$\frac{|1-\lambda|}{|1+\lambda|} = 1. \quad 2$$